

HOMOLOGY EXPONENTS FOR H -SPACES

ALAIN CLÉMENT AND JÉRÔME SCHERER

ABSTRACT. We say that a space X admits a *homology exponent* if there exists an exponent for the torsion subgroup of $H^*(X; \mathbb{Z})$. Our main result states if an H -space of finite type admits a homology exponent, then either it is, up to 2-completion, a product of spaces of the form $B\mathbb{Z}/2^r$, S^1 , $\mathbb{C}P^\infty$, and $K(\mathbb{Z}, 3)$, or it has infinitely many non-trivial homotopy groups and k -invariants. We then show with the same methods that simply connected H -spaces whose mod 2 cohomology is finitely generated as an algebra over the Steenrod algebra do not have homology exponents, except products of mod 2 finite H -spaces with copies of $\mathbb{C}P^\infty$ and $K(\mathbb{Z}, 3)$.

INTRODUCTION

The study of the torsion in the homotopy groups and the integral homology groups of a space motivated the Moore conjecture, see [Sel88], and the Serre conjecture, [Ser53]. Serre proved that a simply connected space with finite dimensional (and non-trivial) mod p (co)homology $H^*(X; \mathbb{F}_p)$ must have infinitely many non-trivial homotopy groups. He conjectured that there should in fact exist infinitely many homotopy groups of X containing p -torsion, which was proved eventually by McGibbon and Neisendorfer [MN84], relying on Miller's solution [Mil84] of the Sullivan conjecture. This was then refined further by Lannes and Schwartz in [LS86]. Their criterion is that $H^*(X; \mathbb{F}_p)$ is locally finite, as a module over the Steenrod algebra. Dwyer and Wilkerson went one step further, [DW90], looking only at the module $QH^*(X; \mathbb{F}_p)$ of indecomposable elements. Félix, Halperin, Lemaire, and Thomas provided yet another criterion involving the depth of $H_*(\Omega X; \mathbb{F}_p)$, [FHLT89]. In their subsequent paper [FHT92] they focused on the size of the torsion part in the “loop space homology” $H_*(\Omega X; \mathbb{Z})$. They proved in fact a homological version of the Moore conjecture, namely that the p -torsion part of the integral homology of the loop space of a $\mathbb{Z}_{(p)}$ -elliptic space always has an exponent.

In this article we are interested in understanding when the torsion subgroup of the integral homology of a large class of loop spaces, and more generally H -spaces, can have an exponent. In the spirit of Serre's theorem, we first classify those H -spaces having a homology exponent at the prime 2 which are Postnikov pieces (they have only a finite number of non-trivial homotopy groups). Thus we will say henceforth that a space admits a *homology exponent* if there exists an integer k such that $2^k \cdot T_2 H^*(X; \mathbb{Z}) = 0$, where T_2 stands for the 2-torsion subgroup. We work with connected H -spaces of finite type.

Theorem 5.2 *Let X be an H -space of finite type which admits a homology exponent. Then either X is, up to 2-completion, a product of spaces of the form $B\mathbb{Z}/2^r$, S^1 , $\mathbb{C}P^\infty$ and $K(\mathbb{Z}, 3)$, or X admits infinitely many non-trivial k -invariants and homotopy groups.*

The methods we develop predict in fact explicit degrees in which to find homology classes of order 2^r for arbitrarily large r when the space has no homology exponent, quite in the spirit of Browder's “infinite implications”, [Bro61]. This builds on previous work by the first author, who analyzed the case of a Postnikov piece with at most two non-trivial homotopy groups in [Clé06].

2000 *Mathematics Subject Classification.* Primary 57T25, 55S45 ; Secondary 55P20, 55S10, 55T10, 55T20.

The second author is supported by the program Ramón y Cajal, MEC, Spain, and FEDER/MEC grant MTM2004-06686. This research was partially supported by the Swiss National Science Foundation grant FN 200020-105383.

There is a class of H -spaces which is very close to the Postnikov pieces we have been dealing with up to now, namely those H -spaces for which the mod 2 cohomology is finitely generated as an algebra over the Steenrod algebra. They are obtained indeed as extensions by H -fibrations of an H -space with finite mod 2 cohomology by a Postnikov piece, [CCSa].

Theorem 7.5 *Let X be a simply connected H -space of finite type such that $H^*(X; \mathbb{F}_2)$ is finitely generated as an algebra over the Steenrod algebra. Assume that X admits a homology exponent. Then X is, up to 2-completion, the product of a mod 2 finite H -space Y with copies of $K(\mathbb{Z}, 2)$ and $K(\mathbb{Z}, 3)$.*

This contrasts with the homological version of the Moore conjecture obtained by Félix, Halperin, and Thomas in [FHT92]. Of course the mod 2 cohomology of the loop space on a finite complex is very rarely finitely generated as an algebra over the Steenrod algebra.

Acknowledgements. We would like to thank Richard Kane for providing a simple proof of Lemma 3.3, and Juan A. Crespo and Wolfgang Pitsch for helpful comments. The second author would like to thank Kathryn Hess and the IGAT, EPFL, for the invitation which made this collaboration possible.

1. REDUCTION TO SIMPLY CONNECTED SPACES

In this section we explain how to reduce the study of arbitrary connected H -spaces to simply connected ones for which the second homotopy group is torsion. These are then the spaces we study in the rest of the article. Let us start with basic terminology and notation.

Notation 1.1. A space X is a *Postnikov piece* if it has only finitely many non-trivial homotopy groups. It is an H -Postnikov piece if it is moreover an H -space. The n -th *Postnikov section* $i_n : X \rightarrow X[n]$ is determined, up to homotopy, by the property that it induces isomorphisms on homotopy groups π_i , for $i \leq n$, and $\pi_i X[n] = 0$ for $i > n$. The homotopy fiber $X\langle n \rangle$ of i_n is the n -connected cover of X . When X is simple (for example when X is simply connected or when X is an H -space), there exist k -invariants $k_n \in H^{n+1}(X[n-1]; \pi_n X)$ such that $X[n]$ can be recovered as the homotopy fiber of a map $k_n : X[n-1] \rightarrow K(\pi_n X, n+1)$ representing the k -invariant. When X is an H -space, all k -invariants are primitive elements.

Let X be a space. By $\{B_r^*, d_r\}$ we denote its mod-2 cohomology *Bockstein spectral sequence*: $B_1^* \cong H^*(X; \mathbb{F}_2) \implies (H^*(X; \mathbb{Z})/\text{torsion}) \otimes \mathbb{F}_2$. Recall that the first differential $d_1 = Sq^1$ is the Bockstein and a pair of elements x and y which survive to the page B_r and such that $d_r(x) = y$ detect a copy of $\mathbb{Z}/2^r$ in $H^*(X; \mathbb{Z})$ in degree $|y| = |x| + 1$.

We collect now a result about “small” Postnikov pieces. These will turn out to be the only H -Postnikov pieces having an exponent.

Proposition 1.2. *Let X be a connected H -space of finite type such that $\pi_2(X)$ and $\pi_3(X)$ are torsion free. Then $X[3]$ is a product of spaces of the form $B\mathbb{Z}/p^r$, S^1 , $\mathbb{C}P^\infty$ and $K(\mathbb{Z}, 3)$. Moreover, $X[3]$ admits a homology exponent.*

Proof. It is well-known that a copy of the integers in $\pi_1 X$ corresponds to a copy of S^1 splitting of X (because of the existence of a section $S^1 \rightarrow X$). One readily verifies that $H^3(K(\mathbb{Z}/p^r, 1); \mathbb{Z}) = 0$, which shows that the first k -invariant must be trivial. Thus $X[2]$ splits as a product of copies of $B\mathbb{Z}/p^r$'s, S^1 's, and $\mathbb{C}P^\infty$'s. Next, the only elements in $H^4(K(\mathbb{Z}/p^r, 1); \mathbb{Z})$, for any prime p and any integer r , and $H^4(K(\mathbb{Z}, 1); \mathbb{Z})$ are multiples of the square of the generator in degree 2. Such elements are not primitive (unless they are trivial), and hence cannot be the k -invariants of an H -space. Therefore, the second k -invariant of $X[3]$ is trivial as well and the space splits as a product.

It remains to prove the assertion about the homology exponent. Recall that $p^r \cdot \tilde{H}^*(B\mathbb{Z}/p^r; \mathbb{Z}) = 0$ (by a transfer argument), $H^*(S^1; \mathbb{Z})$, $H^*(\mathbb{C}P^\infty; \mathbb{Z})$ are torsion free and $2 \cdot T_2(H^*(K(\mathbb{Z}, 3); \mathbb{Z})) = 0$ (as a consequence of Serre [Ser53] or Cartan's computations [Car55]). $\square$

Lemma 1.3. *Let $r \geq 1$ and $(S^1)^r \rightarrow Y \rightarrow X$ be a H -fibration. If X admits a homology exponent, then so will Y .*

Proof. Let us remark that the fibration is orientable [Spa81, p. 476]. We obtain the result for $r = 1$ by inspecting the associated Gysin cohomology exact sequence and conclude by induction on r . $\square$

We conclude this section with the promised reduction. The existence of a homology exponent for an arbitrary H -space is detected in the homology of a certain covering space.

Proposition 1.4. *Let X be a connected H -space of finite type. Then there exists a simply connected H -space of finite type Y such that $\pi_2 Y$ is a torsion abelian group and Y fits into the following H -fibration:*

$$Y \longrightarrow X \longrightarrow B\pi_1 X \times K(\mathbb{Z}^k, 2),$$

for some $k \geq 0$. Moreover, if X admits a homology exponent, then so will Y .

Proof. Let us first deal with the copies of $\mathbb{Z}$ in $\pi_1 X$. They correspond to a torus $(S^1)^r$ splitting off X . There exists hence X_1 such that $X \simeq X_1 \times (S^1)^r$ and $\pi_1 X_1 \cong \bigoplus_a \mathbb{Z}/2^{s_a} \oplus A$, where A is a $2'$ -torsion abelian group. For any a there is a map $B\mathbb{Z}/2^{s_a} \rightarrow K(\mathbb{Z}, 2)$ corresponding to the Bockstein operation of order s_a . Let us define X_2 to be the homotopy fiber of the composite map

$$X_1 \rightarrow B\pi_1 X_1 \rightarrow \prod_a K(\mathbb{Z}, 2) \times K(A, 1).$$

As in the proof of [LS86, Proposition 0.7], X_2 splits as a product $\tilde{X} \times \prod_a S^1$, where $\tilde{X}$ denotes the universal cover of X . By the previous lemma, X_2 , and thus $\tilde{X}$, admit an exponent if X does.

Finally let us write $\pi_2 X \cong \mathbb{Z}^k \oplus A'$ where A' is a finite torsion group and define Y to be the homotopy fiber of the map $\tilde{X} \rightarrow K(\mathbb{Z}^k, 2)$ so that $\pi_2 Y \cong A'$. The previous lemma yields the statement about the exponent. The description of the base of the H -fibration comes from Proposition 1.2. $\square$

2. A SPLITTING PRINCIPLE

Let X be a Postnikov piece, which highest non-trivial homotopy group is $\pi_n X = H$. In this section we show that, if H is torsion free, the $(n-2)$ -connected cover $X(n-2)$ splits as a product $K(H, n) \times K(G, n-1)$. Loosely speaking, the n -th k -invariant attaches $K(H, n)$ directly to $X[n-2]$ and cannot tie the last two homotopy groups together.

We will first need some basic results on Eilenberg-MacLane spaces. We follow the terminology and notation of [Sch94, Chapter 1]. For a unified treatment of the spaces $K(\mathbb{Z}/2^s, n)$, with $s \geq 1$, and $K(\mathbb{Z}, n)$, it is convenient to introduce a notation for the higher Bockstein operations. Let u_n (respectively $Sq_s^1 u_n$) denote the generator of the 1-dimensional $\mathbb{F}_2$ -vector space $H^n(K(\mathbb{Z}/2^s, n); \mathbb{F}_2)$ (respectively $H^{n+1}(K(\mathbb{Z}/2^s, n); \mathbb{F}_2)$). For an admissible sequence $I = (i_1, \dots, i_m)$, we will write $Sq_s^I u_n$ instead of $Sq^{(i_1, \dots, i_{m-1})} Sq_s^1 u_n$ if $i_m = 1$ and instead of $Sq^I u_n$ if $i_m \neq 1$. We denote also by u_n the generator of $H^n(K(\mathbb{Z}, n); \mathbb{F}_2)$.

Serre computed the mod-2 cohomology of Eilenberg-MacLane spaces.

Theorem 2.1. (Serre, [Ser53]) *Let $n \geq 1$ and $s \geq 1$.*

- (1) *The $\mathbb{F}_2$ -algebra $H^*(K(\mathbb{Z}/2^s, n); \mathbb{F}_2)$ is isomorphic to the polynomial $\mathbb{F}_2$ -algebra on generators $Sq_s^I u_n$, where I covers all the admissible sequences of excess $e(I) < n$.*

- (2) The $\mathbb{F}_2$ -algebra $H^*(K(\mathbb{Z}, n); \mathbb{F}_2)$ is isomorphic to the polynomial $\mathbb{F}_2$ -algebra on generators $Sq^I u_n$, where I covers all the admissible sequences of the form $(i_1, \dots, i_n)$ where $i_n \neq 1$ and of excess $e(I) < n$. $\square$

Our first lemma relies on Serre's computations in low degrees.

Lemma 2.2. *Let G be a finitely generated abelian group, H be free abelian, and $n \geq 3$. Then $P^{n+1}H^*(K(G, n-1); H) = 0$.*

Proof. Since $H^*(K(G, n-1); H) \cong H^*(K(G, n-1); \mathbb{Z}) \otimes H$, it is enough to consider the case when $H = \mathbb{Z}$. When $n \geq 4$, the only elements in $H^{n+1}(K(G, n-1); \mathbb{F}_2)$ are sums of elements of the form $Sq^2 u_{n-1}$. These elements all have non-trivial Bockstein and we see therefore from the Bockstein spectral sequence that $H^{n+1}(K(G, n-1); \mathbb{Z})$ is 2-torsion free. As there is obviously no odd primary torsion in this degree (from Cartan's description of $H^*(K(G, n-1); \mathbb{F}_p)$, [Car55]), we have that $H^{n+1}(K(G, n-1); \mathbb{Z}) = 0$. When $n = 3$, write $G \cong \mathbb{Z}^k \oplus A$, where A is torsion. Then $H^4(K(G, 2); \mathbb{Z}) \cong H^4(K(\mathbb{Z}^k, 2); \mathbb{Z})$. There are no primitive elements in this degree. $\square$

Proposition 2.3. *Let X be a simply connected H -space of finite type such that $\pi_2 X$ is a torsion group. Let $n \geq 3$, consider the Postnikov section $X[n]$ and assume that $\pi_n X$ is torsion free. Then we have the following H -fibration:*

$$X[n] \longrightarrow X[n-2] \longrightarrow K(\pi_{n-1} X, n) \times K(\pi_n X, n+1).$$

Proof. Set $G = \pi_{n-1} X$. We prove that the k -invariant $k_n \in H^{n+1}(X[n-1]; H)$ factors through $H^{n+1}(X[n-2]; H)$. Let us consider the fibration

$$K(G, n-1) \rightarrow X[n-1] \rightarrow X[n-2]$$

and the cofibration $K(G, n-1) \rightarrow X[n-1] \rightarrow C$. From Lemma 2.2 we deduce that the H -map $K(G, n-1) \rightarrow X[n-1] \xrightarrow{k_n} K(H, n+1)$ is null-homotopic.

Therefore k_n factors through a map $C \rightarrow K(H, n+1)$. By Ganea's result [Gan65] the fiber of the map $C \rightarrow X[n-2]$ is the join $K(G, n-1) * \Omega(X[n-2])$. This is an n -connected space and $\pi_{n+1}(K(G, n-1) * \Omega(X[n-2])) \cong G \otimes \pi_2 X$, a torsion group because $\pi_2 X$ is so. Thus $K(H, n+1)$ is a $K(G, n-1) * \Omega(X[n-2])$ -local space, as we assume that H is torsion free. From Dwyer's version of Zabrodsky lemma [Dwy96, Proposition 3.5] we deduce that k_n factors through a map $k : X[n-2] \rightarrow K(H, n+1)$.

If $k_{n-1} \in H^n(X[n-2]; G)$ denotes the previous k -invariant, this means that $X[n]$ is the homotopy fiber of the product map $X[n-2] \xrightarrow{k_{n-1} \times k} K(G, n) \times K(H, n+1)$. $\square$

3. GAPS IN THE PRIMITIVES

This section contains the key cohomological result which makes the analysis of the Serre spectral sequence possible. We notice first that there are gaps in the mod 2 cohomology of Eilenberg-MacLane spaces and show then that these gaps propagate in the cohomology of any Postnikov piece.

Definition 3.1. Let $n \geq 1$. We set $A_n = \{a \in \mathbb{N}, a \text{ odd} \mid \nu_2(a) \geq n+1\}$ where $\nu_2(a)$ is the 2-adic length of the integer a .

We will show that there are no indecomposable elements in the cohomology of an n -stage Postnikov piece in degrees $a \in A_n$. To deduce that there are no primitive elements either, we make use of the relationship provided by the Milnor-Moore theorem [MM65, Proposition 4.21]: For a connected, associative, and commutative Hopf algebra over $\mathbb{F}_2$, there is an exact sequence of graded modules

$$0 \longrightarrow P(\xi H) \longrightarrow PH \longrightarrow QH,$$

where ξH is the image of the *Frobenius map* $\xi : x \mapsto x^2$, QH is the module of indecomposable elements and PH is the module of primitive elements of H .

Lemma 3.2. *Let H be a finitely generated abelian group and $n \geq 2$. Then*

$$Q^a H^*(K(H, n); \mathbb{F}_2) = 0 = P^a H^*(K(H, n); \mathbb{F}_2)$$

for all $a \in A_n$.

Proof. When it is not trivial, the $\mathbb{F}_2$ -algebra structure of $H^*(K(H, n); \mathbb{F}_2)$ is given by a polynomial algebra on generators of the form $Sq_s^I u_n$ where I runs over admissible sequences with excess $e(I) < n$, as we have recalled in Theorem 2.1. Careful calculations show that these generators lie in degrees $1 + 2^{h_1} + \dots + 2^{h_{n-1}}$ where $h_1 \geq \dots \geq h_{n-1} \geq 0$ (see [Ser53, Théorème 1, p. 212 and Théorème 2, p. 213]). The 2-adic length of these degrees is bounded by n . This shows that there are no indecomposable elements in the degrees we claimed. These degrees being odd, there are no primitives either, because the kernel of the map $PH^*(K(H, n); \mathbb{F}_2) \rightarrow QH^*(K(H, n); \mathbb{F}_2)$ is concentrated in even degrees. $\square$

The proof of the following lemma has been kindly communicated to us by Richard Kane, [Kan].

Lemma 3.3. *Let B be a connected, associative, and commutative Hopf algebra of finite type over $\mathbb{F}_2$ and A a sub-Hopf algebra of B . Then the morphism $QA \rightarrow QB$ is injective in odd degrees.*

Proof. We work in degree $2n + 1$. Consider the Hopf subalgebra C of A , and hence of B , generated by the elements in A of degree $\leq 2n$. Then one has an inclusion of quotient Hopf algebras $A//C \hookrightarrow B//C$ by [Kan88, Corollary p.9]. Let x be an indecomposable element in QA of degree $2n + 1$. It determines a non-zero primitive element in $P(A//C)$, hence in $P(B//C)$. As the map $P(B//C) \rightarrow Q(B//C)$ is injective in odd degrees, we see that the composite $QA \rightarrow Q(B//C)$ is injective in degree $2n + 1$. Therefore $QA \rightarrow QB$ must be injective in degree $2n + 1$ as well. $\square$

Remark 3.4. The preceding lemma has a nice interpretation in terms of André-Quillen homology, the derived functor of $Q(-)$. It is proved in [CCSb, Proposition 1.3] that one has, in the setting of the lemma, an exact sequence $H_1^Q(B//A) \rightarrow QA \rightarrow QB \rightarrow Q(B//A) \rightarrow 0$, a result dual to that of Bousfield, [Bou70, Theorem 3.6]. Moreover the graded $\mathbb{F}_2$ -vector space $H_1^Q(B//A)$ is concentrated in even degrees.

We are now ready to prove that the gaps also appear in the cohomology of any Postnikov piece.

Proposition 3.5. *Let $n \geq 2$ and X be a simply connected n -stage H -Postnikov piece of finite type. Then $Q^a H^*(X; \mathbb{F}_2) = 0 = P^a H^*(X; \mathbb{F}_2)$ for all $a \in A_n$.*

Proof. The proof goes by induction on n . We have the following H -fibration given by the Postnikov tower of X :

$$K(\pi_n(X), n) \longrightarrow X \xrightarrow{p} X[n-1].$$

We rely on the analysis of the Eilenberg-Moore spectral sequence done by Smith [Smi70, Proposition 3.2]. The quotient Hopf algebra $R = H^*(X[n-1]; \mathbb{F}_2) // \ker p^*$ can be identified via p^* with a sub-Hopf algebra of $H^*(X; \mathbb{F}_2)$. The corresponding quotient $S = H^*(X; \mathbb{F}_2) // R$ is isomorphic to a sub-Hopf algebra (and a sub- $\mathcal{A}_2$ -algebra) of $H = H^*(K(\pi_n(X), n); \mathbb{F}_2)$. There is a section $S \rightarrow H^*(X; \mathbb{F}_2)$, which is a map of algebras, so that the module of indecomposables $QH^*(X; \mathbb{F}_2)$ is isomorphic to $QR \oplus QS$, as graded $\mathbb{F}_2$ -vector spaces. We have to prove that both QR and QS are trivial in degrees in A_n . First, since $Q(-)$ is right exact, we have a surjection $QH^*(X[n-1]; \mathbb{F}_2) \twoheadrightarrow QR$. Now, $Q^a H^*(X[n-1]; \mathbb{F}_2) = 0$ for any $a \in A_{n-1}$ by induction hypothesis and we conclude that $Q^a R = 0$ for any $a \in A_n$ since $A_n \subset A_{n-1}$. Second, we deal with QS . Let us apply the preceding lemma to the inclusion $S \subset H^*(X; \mathbb{F}_2)$. We see that $QS \rightarrow QH$ is a monomorphism in odd degrees. Therefore $Q^a S = 0$ for all $a \in A_n$ by Lemma 3.2. $\square$

4. TRANSVERSE ELEMENTS IN EILENBERG-MAC LANE SPACES

Now begins the study of the 2-torsion in Postnikov pieces. In this section we deal with the first step of the induction, namely the analysis of the case of Eilenberg-MacLane spaces. Recall that $\{B_r^*, d_r\}$ denote the mod-2 cohomology Bockstein spectral sequence of a space X .

Definition 4.1. Let n and r be two positive integers. An element $x \in B_r^n$ is said to be ℓ -transverse if $d_{r+l}x^{2^l} \neq 0 \in B_{r+l}^{2^l n}$ for all $0 \leq l \leq \ell$. An element $x \in B_r^n$ is said to be *transverse* if it is ℓ -transverse for all $\ell \geq 0$. We will also speak of *transverse implications* of an element $x \in B_r^n$.

Every transverse element gives rise to 2-torsion of arbitrarily high order in the integral cohomology of X . This definition, introduced in [Clé02], adapts Browder's "infinite implications" from [Bro61] to our purpose. To us, the fact that the elements die in increasing pages of the Bockstein spectral sequence is crucial, whereas Browder was merely interested to know that the degrees of the elements was increasing.

Our strategy for disproving the existence of a homology exponent for a space will consist in exhibiting a transverse element in its mod-2 cohomology Bockstein spectral sequence. Note that in principle the absence of transverse elements does not imply the existence of an exponent for the 2-torsion part in $H^*(X; \mathbb{Z})$. An easy example is given by the infinite wedge $\vee M(\mathbb{Z}/2^n, n)$.

In the special case of Eilenberg-MacLane spaces, we have the following result, taken from the first author PhD thesis, [Clé02, Theorem 1.3.2].

Theorem 4.2. *Let H be an abelian group of finite type and let $n \geq 2$. Consider the Eilenberg-MacLane space $K(H, n)$ and its mod-2 cohomology Bockstein spectral sequence $\{B_r^*, d_r\}$. Suppose that one of the following assumptions holds:*

- n is even and $x \in B_{s_j}^n$ is 0-transverse for some $1 \leq j \leq l$,
- $x \in P^{\text{even}} B_1^*$ is 0-transverse ($Sq^1 x \neq 0$).

Then x is transverse. □

Note that the abelian group H is isomorphic to $\mathbb{Z}^s \oplus \mathbb{Z}/2^{s_1} \oplus \cdots \oplus \mathbb{Z}/2^{s_l} \oplus A$, where A is a $2'$ -torsion group, which is therefore invisible to the mod 2 Bockstein spectral sequence. Hence the first type of 0-transverse elements correspond basically to the fundamental classes u_n introduced in Section 2, one for each copy of $\mathbb{Z}/2^{s_j}$ (the fundamental classes coming from the copies of $\mathbb{Z}$ survive to B_∞^n).

Remark 4.3. In general a 0-transverse implication does not imply transverse implications. More precisely, the fact that $x \in P^{\text{even}} H^*(X; \mathbb{F}_2)$ is such that $Sq^1 x \neq 0$ does not always force x to be transverse. A counter-example is given by $X = BSO$ and $x = w_2$, the second Stiefel-Whitney class in $H^2(BSO; \mathbb{F}_2)$.

From Theorem 4.2 it is not difficult to prove that most Eilenberg-MacLane spaces have no homology exponent.

Proposition 4.4. *Let H be a non-trivial 2-torsion abelian group and let $n \geq 2$. The Eilenberg-MacLane space $K(H, n)$ has no homology exponent.*

Proof. Accordingly to the Künneth formula, it is sufficient to establish the result when $H = \mathbb{Z}/2^s$ for some $s \geq 1$. If n is even, consider the reduction of the fundamental class $u_n \in H^n(K(\mathbb{Z}/2^s, n); \mathbb{F}_2)$. This class survives to B_s^n and is 0-transverse. Then $u_n \in B_s^n$ is transverse. If n is odd, consider the admissible sequence $(2, 1)$. Its excess is exactly 1 and therefore $Sq_s^{2,1} u_n \in P^{\text{even}} H^*(K(\mathbb{Z}/2^s, n); \mathbb{F}_2)$ when $n \geq 3$. Moreover we have $Sq^1 Sq_s^{2,1} u_n = Sq_s^{3,1} u_n$ by Adem relations, which means that $Sq_s^{2,1} u_n$ is 0-transverse. Hence $Sq_s^{2,1} u_n \in B_1^{n+3}$ is transverse. □

Proposition 4.5. *Let H be a finitely generated abelian group and $n \geq 4$. The Eilenberg-MacLane space $K(H, n)$ is then either mod 2 acyclic, or has no homology exponent.*

Proof. By the Künneth formula and Proposition 4.4, it is sufficient to analyze the case $H = \mathbb{Z}$. Consider the reduction of the fundamental class $u_n \in H^n(K(\mathbb{Z}, n); \mathbb{F}_2)$. If n is even, then $Sq^2 u_n$ is transverse. If n is odd, then $Sq^{6,3} u_n$ is transverse. $\square$

5. TRANSVERSE ELEMENTS IN POSTNIKOV PIECES

We are now ready to prove our main result: Most Postnikov pieces do not have a homology exponent. The strategy to prove this relies on the crucial observation that the transverse implications of certain element in the cohomology of the total space of a fibration can be read in the cohomology of the fibre.

Lemma 5.1. *Let $j : F \rightarrow X$ be a continuous map. If $x \in H^*(X; \mathbb{F}_2)$ is such that $j^*(x) \neq 0 \in H^*(F; \mathbb{F}_2)$ is transverse, then x itself is transverse.*

Proof. It follows from the naturality of the Bockstein spectral sequence. $\square$

Theorem 5.2. *Let X be an H -space of finite type which admits a homology exponent. Then either X is, up to 2-completion, a product of spaces of the form $B\mathbb{Z}/2^r$, S^1 , $\mathbb{C}P^\infty$ and $K(\mathbb{Z}, 3)$, or X admits infinitely many non-trivial k -invariants and homotopy groups.*

Proof. Let us assume that X is a Postnikov piece. By Proposition 1.4, there is an H -fibration of the form

$$Y \longrightarrow X \longrightarrow B\pi_1 X \times K(\mathbb{Z}^r, 2),$$

where Y is a simply connected H -space of finite type such that $\pi_2 Y$ is a torsion abelian group. Moreover, Y admits a homology exponent. It is also clearly a Postnikov piece. Let us show that Y is, up to 2-completion, a product of copies of $K(\mathbb{Z}, 3)$. By Proposition 1.2, this will imply that X itself splits as the announced product.

Assume that $\pi_n Y = H$ is the highest non-trivial homotopy group of Y , up to 2-completion. If $n = 2$, since $\pi_2 Y$ is a torsion abelian group, we deduce from Proposition 4.4 that H is $2'$ -torsion. In other words $Y_2^\wedge$ is contractible. We can therefore assume that $n \geq 3$. The space Y fits into the fibration sequence

$$K(H, n) \xrightarrow{j} Y \xrightarrow{i} Y[n-1] \xrightarrow{k} K(H, n+1),$$

where k denotes the last k -invariant. We analyze the situation in two steps, depending on the presence of 2-torsion in H .

Let us first assume that H contains 2-torsion, let us say $\oplus_b \mathbb{Z}/2^{t_b}$. Choose an index b and consider the projection $\pi : H \rightarrow \mathbb{Z}/2^{t_b}$ on the corresponding cyclic subgroup. Pick $v_n \in H^n(K(H, n); \mathbb{F}_2)$, the image via π^* of the class $u_n \in H^n(K(\mathbb{Z}/2^{t_b}, n); \mathbb{F}_2)$.

Set $\xi = (2^{n-1} - 2, 2^{n-2} - 1, 2^{n-3} - 1, \dots, 3, 1)$. The degree $\deg(Sq_t^\xi v_n) = 2^n - 2$ is even and $Sq^1 Sq_t^\xi v_n \neq 0$ since $e(\xi) = n - 2$. By Theorem 4.2, $Sq_t^\xi v_n$ is transverse. Since Y is an H -space and the k -invariant is an H -map, the element $d_{n+1} v_n$ is primitive, and so is $d_{2^n-1} Sq_t^\xi v_n = Sq_t^\xi d_{n+1} v_n \in P^{2^n-1} H^*(Y[n-1]; \mathbb{F}_2)$. By Proposition 3.5, $P^{2^n-1} H^*(Y[n-1]; \mathbb{F}_2) = 0$ since $2^n - 1 \in A_{n-1}$. Therefore, $Sq_t^\xi v_n$ survives in the Serre spectral sequence and by the previous lemma, $H^*(Y; \mathbb{F}_2)$ contains a transverse element. In particular it has no homology exponent.

Hence, H must be 2-torsion free and is thus isomorphic to $\mathbb{Z}^s \oplus A$, where A is a torsion group, for some $s \geq 1$. By Proposition 2.3, Y fits in the following H -fibration:

$$K(H, n) \times K(\pi_{n-1} Y, n-1) \longrightarrow Y \longrightarrow Y[n-2].$$

Choose now $v_n \in H^n(K(H, n); \mathbb{F}_2)$ to be the image of the class $u_n \in H^n(K(\mathbb{Z}, n); \mathbb{F}_2)$ given by projection on the first copy of $\mathbb{Z}$ in H .

If $n \geq 4$, then set $\eta = (2^{n-2} + 2^{n-3} - 2, 2^{n-3} + 2^{n-4} - 1, 2^{n-4} + 2^{n-5} - 1, \dots, 5, 2)$. The degree $\deg(Sq^\eta v_n) = 2^{n-1} + 2^{n-2} - 2$ is even and $Sq^1 Sq^\eta v_n \neq 0$ since $e(\eta) = n - 2$. Thus $Sq^\eta v_n$ is transverse

and survives in the Serre spectral sequence of the above fibration since $2^{n-1} + 2^{n-2} - 1 \in A_{n-2}$. In this case, $H^*(Y; \mathbb{F}_2)$ contains a transverse element and has no homology exponent.

Therefore, $n = 3$ and $Y \simeq K(H, 3) \times K(\pi_2 Y, 2)$. Since Y admits a homology exponent, the torsion group $\pi_2 Y$ is trivial and H is torsion free. $\square$

The proof of the theorem predicts explicit degrees in which to find higher and higher torsion in the integral cohomology of the space.

Corollary 5.3. *Let X be a simply connected H -Postnikov piece of finite type, say $X \simeq X[n]$. Assume that $\pi_2 X$ is torsion and that X is not equivalent up to 2-completion to a product of copies of $K(\mathbb{Z}, 3)$. Then, for any integer k , there is a copy of $\mathbb{Z}/2^k$ in $H^*(X; \mathbb{Z})$*

- (1) *in degree $2^k(2^n - 2)$ if $\pi_n X$ contains 2-torsion,*
- (2) *in degree $2^k(2^{n-1} + 2n - 2 - 2)$ if not.*

Proof. Since X is different from $K(\mathbb{Z}^m, 3)$, we know from Theorem 5.2 that X has no exponent. The higher and higher torsion is detected by the consecutive powers of the elements $Sq^\xi v_n$ and $Sq^\eta v_n$ constructed in the above proof. $\square$

Any finite H -space has obviously a homology exponent. Our second corollary applies to its Postnikov sections. As soon as it has at least two homotopy groups, it cannot have a homology exponent.

Corollary 5.4. *Let X be a simply connected finite H -space and $n \geq 3$. Then $X[n]$ has a homology exponent if and only if $X[n] \simeq X[3] \simeq K(\mathbb{Z}^r, 3)$ for some $r \geq 0$.*

Proof. The fact that the H -space X is finite and simply connected forces it to be 2-connected, [Bro61, Theorem 6.10]. Moreover, $\pi_3 X \cong \mathbb{Z}^r$ for some integer r , by work of Hubbuck and Kane, [HK75]. The result now follows directly from Theorem 5.2. $\square$

This corollary applies in particular to S^3 . The Postnikov section $S^3[3] \simeq K(\mathbb{Z}, 3)$ has a homology exponent, but all higher Postnikov sections $X[n]$, $n \geq 4$, have none. The following proof of a result obtained by Levi in [Lev95] is, to our knowledge, the first one not based on Miller's solution of the Sullivan's conjecture [Mil84]. Let us mention in this context the work of Klaus, [Kla02], who proves the statement about the k -invariants for $BG_2^\wedge$, not for the loop space.

Corollary 5.5. *Let G be a 2-perfect finite group. Then $\Omega(BG_2^\wedge)$ has infinitely many non-trivial k -invariants and homotopy groups.*

Proof. Suppose $BG_2^\wedge$ is a Postnikov piece. Following Levi [Lev95], there is a homology exponent for $\Omega(BG_2^\wedge)$ and therefore this space has to be a product of copies of $B\mathbb{Z}/2^r$, S^1 , CP^∞ or $K(\mathbb{Z}, 3)$. Since $\Omega(BG_2^\wedge)$ has torsion homotopy groups, the only copies that can occur are of the form $B\mathbb{Z}/2^r$. Thus $BG_2^\wedge \simeq K(A, 2)$, where A is a 2-torsion abelian group. By the Evens-Venkov theorem, [Eve61], $H^*(BG_2^\wedge; \mathbb{F}_2)$ is Noetherian. Hence A is trivial, and so is $BG_2^\wedge$. $\square$

6. COMPARISON WITH OTHER FORMS OF SERRE'S THEOREM

In this section we compare our theorem to the other results we mentioned in the introduction. We show that the existence of a homology exponent is stronger than all previously established criteria, except possibly [FHLT89], which seems difficult to relate directly to cohomological statements. Therefore, when X is an H -space, our result provides new proofs of those. They are very different in spirit, since they do not require the Sullivan conjecture. For simplicity we deal here with simply connected spaces.

Proposition 6.1. *Let X be a simply connected H -Postnikov piece. Then*

- (1) (Serre [Ser53]) *$H^*(X; \mathbb{F}_2)$ is not finite,*

- (2) (Lannes-Schwartz [LS86]) $H^*(X; \mathbb{F}_2)$ is not locally finite,
- (3) there exists an element of infinite height in $H^*(X; \mathbb{F}_2)$,
- (4) (Grodal [Gro98]) the transcendence degree of $H^*(X; \mathbb{F}_2)$ is infinite unless X is homotopy equivalent, up to 2-completion, to $K(\mathbb{Z}, 2)^s$,
- (5) (Dwyer-Wilkerson [DW90]) the unstable module $QH^*(X; \mathbb{F}_2)$ is not locally finite unless X is homotopy equivalent, up to 2-completion, to $K(\mathbb{Z}, 2)^s$.

Proof. Notice first that $K(\mathbb{Z}, 2)$ and $K(\mathbb{Z}, 3)$ satisfy (1) - (5). Assume now that X is a Postnikov piece, say $X \simeq X[n]$. In the proof of Theorem 5.2 we first considered the covering fibration $(S^1)^r \rightarrow Y \rightarrow X$. The map $Y \rightarrow X$ induces isomorphisms in homology in high degrees. We can therefore assume that $\pi_2 X$ is torsion. Our proof then provides a transverse element $x \in H^*(X; \mathbb{F}_2)$ in even degree whose image in $H^*(K(\pi_n X, n))$ is a transverse element of the form $Sq_t^I u_n$ for some admissible sequence $(i_1, \dots, i_m)$. In particular all powers x^{2^k} are non-zero, which proves (1) - (3). Moreover the elements $x, Sq^{2i_1} x, Sq^{4i_1, 2i_1} x, \dots$ are non-zero, indecomposable, and algebraically independent because so are the corresponding images in $H^*(K(\pi_n X, n))$. This proves (4) and (5). $\square$

7. COHOMOLOGICAL FINITENESS CONDITIONS

The strategy we followed to analyze the integral homology of Postnikov pieces can be applied in a more general context. We work in this last section with simply connected H -spaces X such that $H^*(X; \mathbb{F}_2)$ is finitely generated as an algebra over the Steenrod algebra. This section relies on the Sullivan conjecture. As it may be considered thus as less elementary than the part about Postnikov pieces, we have decided to postpone it till the end of the article.

From the assumption on the mod 2 cohomology, we infer by [CCSa, Lemma 7.1] that there exists an integer n such that the module $QH^*(X; \mathbb{F}_2)$ of indecomposable elements lies in the $(n-1)$ -st stage of the Krull filtration for unstable modules, [Sch94]. Therefore there exists by [CCSa, Theorem 7.3] a simply connected H -space $Y = P_{B\mathbb{Z}/2} X$ with finite mod 2 cohomology and a series of principal H -fibrations

$$X = X_n \xrightarrow{p_n} X_{n-1} \rightarrow \dots \rightarrow X_1 \xrightarrow{p_1} X_0 = Y$$

of simply connected spaces such that the homotopy fiber of p_i is an Eilenberg-Mac Lane space $K(P_i, i)$, where P_i splits as a product of a finite direct sum P_i' of cyclic groups $\mathbb{Z}/2^r$ and a finite direct sum P_i'' of Prüfer groups $\mathbb{Z}_{2^\infty}$. Let us recall here that X_k is obtained as the $\Sigma^k B\mathbb{Z}/p$ -nullification of X (the above tower is Bousfield's nullification tower, [Bou94]). Since $K(\mathbb{Z}_{2^\infty}, i)$ and $K(\mathbb{Z}, i+1)$ are mod 2 equivalent, we alter slightly the way in which the P_i 's are added to Y in order to work in a more familiar setting. Then we can recover X from the tower

$$X = Y_n \xrightarrow{q_n} Y_{n-1} \rightarrow \dots \rightarrow Y_1 \xrightarrow{q_1} Y_0 = Y$$

of simply connected spaces and principal H -fibrations, where the homotopy fiber of q_i is the product of Eilenberg-Mac Lane spaces $K(P'_{i+1}, i+1) \times K(P''_i, i)$. Notice that $Q_1 = P'_1$ is trivial because we assume that X is simply connected ($Y_2^\wedge$ is therefore 2-connected, [Bro61]). We have a splitting result, just like in Proposition 2.3.

Lemma 7.1. *Let X be a simply connected H -space such that $H^*(X; \mathbb{F}_2)$ is finitely generated as an algebra over the Steenrod algebra. Assume that $\pi_2 X_2^\wedge$ is torsion. Then there is an H -fibration*

$$X \longrightarrow Y_{n-2} \longrightarrow K(P''_n \oplus P'_n, n+1) \times K(P''_n, n).$$

Proof. The proof is based on the Zabrodsky lemma, as in Proposition 2.3. $\square$

Our next result is the analog in the present setting of Proposition 3.5. Recall from Section 3 that the set A_n consists in those integers for which the 2-adic length is strictly larger than n .

Proposition 7.2. *Let X be a simply connected H -space such that $H^*(X; \mathbb{F}_2)$ is finitely generated as an algebra over the Steenrod algebra. There exists then integers m and N such that $Q^a H^*(X; \mathbb{F}_2) = 0 = P^a H^*(X; \mathbb{F}_2)$ for all $a \in A_n$ with $a \geq N$.*

Proof. The integer m is determined by the stage of the Krull filtration in which $QH^*(X; \mathbb{F}_2)$ lives, i.e. by the degrees in which the homotopy groups of the homotopy fiber of $X \rightarrow P_{B\mathbb{Z}/2}X = Y$ are non-trivial. With the above notation, $m = n$ if P_n'' is trivial, and $m = n + 1$ if P_n'' is not. The proof goes then by induction on m . When $m = 0$, choose N to be larger than the cohomological dimension of Y . The proof of Proposition 3.5 goes through. $\square$

Lemma 7.3. *Let X be a simply connected H -space which fits, up to 2-completion, in an H -fibration of the form*

$$K(\oplus_t \mathbb{Z}, 2) \longrightarrow X \longrightarrow Y$$

where $H^(Y; \mathbb{F}_2)$ is finite. Then X has no homology exponent unless the fibration splits up to 2-completion, i.e. $X \simeq_2^\wedge Y \times K(\oplus_t \mathbb{Z}, 2)$.*

Proof. Let us omit the 2-completions in the proof and write the details of the proof when $t = 1$. By the result of Hubbuck and Kane, [HK75], $\pi_3 Y$ is isomorphic to a direct sum of say s copies of $\mathbb{Z}$. The map classifying the fibration factors through $Y[3] \simeq K(\oplus_s \mathbb{Z}, 3) \rightarrow K(\mathbb{Z}, 3)$. The E_2 -term of the Serre spectral sequence has the form $\mathbb{Z}[u] \otimes H^*(Y; \mathbb{Z})$, where u has degree 2 and the cohomology of Y is of finite dimension N , and of exponent 2^a for some integer a . The differential $d_3(u) = x$ for some non-zero element $x \in H^3(Y; \mathbb{Z}) \cong \oplus_s \mathbb{Z}$. Therefore $d_3(u^n) = nx \otimes u^{n-1}$. At worst $d_3(x \otimes u^{n-1})$ is non-zero and then hits a torsion element, of order at most 2^a . Hence, on the third column of the E_4 -term, we have a group covering $\mathbb{Z}/2^{n-a}$ in vertical degree 2^n . From the finiteness of Y we see that the spectral sequence collapses at E_{N-3} . An iteration of the above argument shows therefore that the third column of the E_∞ -term contains a group covering $\mathbb{Z}/2^{n-(N-5)a}$ in vertical degree 2^n , for any $n \geq 1$. In particular there is arbitrarily high torsion in $H^*(X; \mathbb{Z})$. Therefore, for X to have an exponent, the fibration must split. $\square$

Remark 7.4. We point out that the preceding lemma provides simple examples of fibrations, such as $K(\mathbb{Z}, 2) \rightarrow S^3 \langle 3 \rangle \rightarrow S^3$, where both the fiber and the base have an exponent, but the total space has none.

Theorem 7.5. *Let X be a simply connected H -space of finite type such that $H^*(X; \mathbb{F}_2)$ is finitely generated as an algebra over the Steenrod algebra. Assume that X admits a homology exponent. Then X is, up to 2-completion, the product of a mod 2 finite H -space Y with copies of $K(\mathbb{Z}, 2)$ and $K(\mathbb{Z}, 3)$.*

Proof. We follow the proof of Theorem 5.2. Let us thus assume that X admits a homology exponent. By killing the copies of $\mathbb{Z}$ in $\pi_2 X$ just like in Proposition 1.4, we can assume that $\pi_2 X_2^\wedge$ is torsion. We also see by inspection of the tower that $\pi_2(Y_i)_2^\wedge$ is torsion for any $i \geq 0$. Therefore the splitting in Lemma 7.1 holds and we work with a fibration

$$K(P_n'' \oplus P_n', n) \times K(P_n'', n-1) \longrightarrow X \longrightarrow Y_{n-2}.$$

If $P_n' \neq 0$, it must contain a copy of $\mathbb{Z}/2^r$ as direct summand. Choose a power of the corresponding element $Sq_t^\xi v_n$, of degree larger than the integer N given in Proposition 7.2. From the Serre spectral sequence for the above fibration we see that this provides a transverse element in $H^*(X; \mathbb{F}_2)$. Therefore $P_n' = 0$ (and so P_n'' is not trivial).

If $n \geq 3$ we choose a copy of $\mathbb{Z}_{2^\infty}$ in P_n'' and a suitable power of the corresponding element $Sq^n v_n$ to detect a transverse element in $H^*(X; \mathbb{F}_2)$. Since we assume that X has a homology exponent, we see that $n \leq 2$, i.e. X is the homotopy fiber of a map $k : Y \rightarrow K(P_1'', 2) \times K(P_2'', 3)$. To conclude the proof we must show that this map is trivial.

The mod 2 cohomology of the H -space Y is finite. Rationally it is thus a product of odd dimensional spheres and, in particular, $\pi_4 Y_2^\wedge$ is torsion. This implies that the projection of k on the second factor $Y \rightarrow K(P_2'', 3)$ is the trivial map, up to 2-completion. Hence the copies of $K(\mathbb{Z}_2^\wedge, 3)$ split off $X_2^\wedge$. We are left with the analysis of a fibration $X \rightarrow Y \rightarrow K(P_1'', 2)$. If the map $Y \rightarrow K(P_1'', 2)$ is not trivial, we conclude from Lemma 7.3 that X cannot have a homology exponent. Hence, the fibration must split and this concludes the proof. $\square$

REFERENCES

- [Bou70] A. K. Bousfield. Nice homology coalgebras. *Trans. Amer. Math. Soc.*, 148:473–489, 1970.
- [Bou94] ———. Localization and periodicity in unstable homotopy theory. *J. Amer. Math. Soc.*, 7: 831–873, 1994.
- [Bro61] W. Browder. Torsion in H -spaces. *Ann. of Math. (2)*, 74:24–51, 1961.
- [Car55] H. Cartan. *Séminaire Henri Cartan de l'Ecole Normale Supérieure, 1954/1955. Algèbres d'Eilenberg-MacLane et homotopie, exposés 2–16*. Secrétariat mathématique, 11 rue Pierre Curie, Paris, 1955.
- [CCSa] N. Castellana, J. A. Crespo, and J. Scherer. Deconstructing Hopf spaces. to appear in *Invent. Math.*
- [CCSb] ———. On the cohomology of highly connected of finite complexes. Preprint, available at: <http://front.math.ucdavis.edu/math.AT/>.
- [Clé02] A. Clément. Integral cohomology of finite Postnikov towers. 2002. Ph. D. Thesis, University of Lausanne, Switzerland.
- [Clé06] ———. Integral cohomology of 2-local Hopf spaces with at most two non-trivial finite homotopy groups. In *An alpine anthology of homotopy theory*, volume 399 of *Contemp. Math.*, pages 87–99. Amer. Math. Soc., Providence, RI, 2006.
- [DW90] W.G. Dwyer and C.W. Wilkerson. Spaces of null homotopic maps. *Astérisque*, (191):6, 97–108, 1990. International Conference on Homotopy Theory (Marseille-Luminy, 1988).
- [Dwy96] W. G. Dwyer. The centralizer decomposition of BG . In *Algebraic topology: new trends in localization and periodicity (Sant Feliu de Guíxols, 1994)*, volume 136 of *Progr. Math.*, pages 167–184. Birkhäuser, Basel, 1996.
- [Eve61] L. Evens. The cohomology ring of a finite group. *Trans. Amer. Math. Soc.*, 101:224–239, 1961.
- [FHLT89] Y. Félix, S. Halperin, J.-M. Lemaire, and J.-C. Thomas. Mod p loop space homology. *Invent. Math.*, 95(2):247–262, 1989.
- [FHT92] Y. Félix, S. Halperin, and J.-C. Thomas. Torsion in loop space homology. *J. Reine Angew. Math.*, 432:77–92, 1992.
- [Gan65] T. Ganea. A generalization of the homology and homotopy suspension. *Comment. Math. Helv.*, 39:295–322, 1965.
- [Gro98] J. Grodal. The transcendence degree of the mod p cohomology of finite Postnikov systems. In *Stable and unstable homotopy (Toronto, ON, 1996)*, volume 19 of *Fields Inst. Commun.*, pages 111–130. Amer. Math. Soc., Providence, RI, 1998.
- [HK75] J. R. Hubbuck and R. Kane. On π_3 of a finite H -space. *Trans. Amer. Math. Soc.*, 213:99–105, 1975.
- [Kan88] R. Kane. *The homology of Hopf spaces*. North-Holland, Amsterdam, 1988.
- [Kan] ———. personal communication. November 2006.
- [Kla02] S. Klaus. A generalization of the non-triviality theorem of Serre. *Proc. Amer. Math. Soc.*, 130(5):1249–1256 (electronic), 2002.
- [Lev95] R. Levi. On finite groups and homotopy theory. *Mem. Amer. Math. Soc.*, 118(567):xiv+100, 1995.
- [LS86] J. Lannes and L. Schwartz. À propos de conjectures de Serre et Sullivan. *Invent. Math.*, 83(3):593–603, 1986.
- [Mil84] H. Miller. The Sullivan conjecture on maps from classifying spaces. *Ann. of Math. (2)*, 120(1):39–87, 1984.
- [MM65] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965.
- [MN84] C. A. McGibbon and J. A. Neisendorfer. On the homotopy groups of a finite-dimensional space. *Comment. Math. Helv.*, 59(2):253–257, 1984.
- [Sch94] L. Schwartz. *Unstable modules over the Steenrod algebra and Sullivan's fixed point set conjecture*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1994.
- [Sel88] P. Selick. Moore conjectures. In *Algebraic topology—rational homotopy (Louvain-la-Neuve, 1986)*, volume 1318 of *Lecture Notes in Math.*, pages 219–227. Springer, Berlin, 1988.
- [Ser53] J.-P. Serre. Cohomologie modulo 2 des complexes d'Eilenberg-MacLane. *Comment. Math. Helv.*, 27:198–232, 1953.
- [Smi70] L. Smith. *Lectures on the Eilenberg-Moore spectral sequence*. Lecture Notes in Mathematics, Vol. 134. Springer-Verlag, Berlin, 1970.
- [Spa81] E. H. Spanier. *Algebraic topology*. Springer-Verlag, New York, 1981. Corrected reprint.

RUE LOUIS-MEYER 9,
CH – 1800 VEVEY,
SWITZERLAND.

E-mail address: `alain.clement@bluewin.ch`

DEPARTAMENT DE MATEMÀTIQUES,
UNIVERSITAT AUTÒNOMA DE BARCELONA,
E-08193 BELLATERRA,
SPAIN.

E-mail address: `jscherer@mat.uab.es`